

# Buschke – Löwenstein Tumor: Presentation of a Case

\*Correspondence:

[paololeonej@gmail.com](mailto:paololeonej@gmail.com)

Address: Av. Pedro J. Menéndez Gilbert y Atahualpa Chávez (junto a la ciudadela Atarazana). Apt. Postal 090505. Departamento de dermatología. Instituto Oncológico Nacional "Dr. Juan Tanca Marengo", Sociedad de Lucha Contra el Cáncer, Solca, Guayaquil, Ecuador. Phone [593] (04) 371 8300 EXT. 2218.

**Conflict of interest:** The authors declare not to have any interest conflicts.

**Received:** June 18; 2022

**Accepted:** July 9; 2022

**Published:** August 19; 2022

**Editor:** Dra. Evelyn Valencia Espinoza

**Cite:**

Leone P, Carvajal M. Buschke – Löwenstein Tumor: Presentation of a Case. Rev. Oncol. Ecu 2022;32(2):253-265.

**DOI:** <https://doi.org/10.33821/619>

Copyright Leone P, et al. This article is distributed under the terms of [Creative Commons Attribution License BY-NC-SA 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows the use and redistribution of citing the source and the original author.

**Paolo Leone Jimenez <sup>1</sup>, Maria Fernanda Carvajal <sup>1</sup>.**

1. Department of Dermatology, National Oncology Institute "Dr. Juan Tanca Marengo," Society for the Fight Against Cancer, Solca, Guayaquil, Ecuador.

## Abstract

**Introduction:** Buschke-Lowenstein tumor (BLT) is a sexually transmitted disease caused by human papillomavirus (HPV), described as an intermediate form between condyloma acuminata and squamous cell carcinoma. It mainly affects the genital and anorectal areas and has the capacity for malignant transformation and a high recurrence rate. Surgery is the first-line treatment.

**Clinical case:** We present the case of a 27-year-old male patient with warty lesions of progressive growth in the inguinal and genital areas. Through the clinical-pathological correlation, the diagnosis of BLT was reached. After discussion in a multidisciplinary committee, it was declared unresectable, and treatment with radiotherapy was resolved, in addition to therapeutic vaccination against HPV, both systemic and intralesional.

**Conclusion:** BLT is locally aggressive and challenging to treat, so prevention against HPV is essential. Therapeutic vaccination in conjunction with radiotherapy offered clinical improvement.

**Keywords:**

MESH: Buschke-Lowenstein Tumor, Condyloma Acuminata, Human Papillomavirus 6, Human Papillomavirus 11, Nonvalent Vaccine, Radiotherapy

DOI: 10.33821/619

## Introduction

Buschke – Löwenstein tumor was first described in 1925 by Abraham Buschke and Ludwig Löwenstein as a neoplasm of the penis that resembled both condyloma acuminata and squamous cell carcinoma [ 1 ]. It is a highly aggressive locoregional tumor, which is why it is said to behave malignantly but with low metastatic activity. It mainly affects the genital area, although rare cases have been described in which it appears in the anorectal area [ 2 ].

Although it is considered histologically benign and is a rare entity, it has a potential for malignant transformation of 8.5–23.8% and reported mortality rates of 20–30% [ 3, 4 ]. The leading cause of this tumor is HPV, generally subtypes 6 and 11, which is considered a sexually transmitted disease [ 5 ].

TBL is a rare entity with no apparent therapeutic guidelines, although surgery is considered the first-line treatment. Next, we present a clinically advanced case where surgery was impossible, making it difficult to manage. Other therapeutic options that could be useful in similar cases were explored.

---

## Clinical case

The patient is a 27-year-old male Hispanic, a man who has sex with men (MSM), who came to the consultation due to multiple warty lesions of 3 years of evolution, which began as small warty plaques on the glans and were treated with podophyllotoxin. Despite treatment, one year before the initial consultation, the lesions recurred and worsened, involving the entire glans, scrotum, and inguinal region.

As a personal history, the patient was HIV+ (human immunodeficiency virus +), diagnosed in September 2019, and was on antiretroviral treatment with lopinavir, ritonavir, atazanavir, tenofovir, and emtricitabine. In addition, he was diagnosed with KS in 2020, for which he received treatment with thalidomide and had a history of chylothorax due to tuberculosis.

The patient provided the report of a biopsy performed in a private center in January 2021 that indicated a viral wart with moderate dysplasia in an unspecified genital region lesion. In the pelvic region, a sizeable cauliflower-like tumor with a warty surface was observed, which compromised the entire scrotum and another that compromised the glans almost wholly, in addition to multiple satellite lesions. This tumor caused limitations in walking and was painful. Bilateral inguinal lymphadenopathies and bilateral edema in the lower extremities were observed and palpated.

## Diagnostic workshop

In the laboratory tests performed, the following was found:

- Erythrocytosis (red blood cells - January 2021:  $5.72 \times 10^6/\mu\text{L}$ . Range:  $4.00 - 5.50 \times 10^6/\mu\text{L}$ )
- Thrombocytosis (Platelets - January 2021:  $584.00 \times 10^3/\mu\text{L}$ ; July 2021:  $466.00 \times 10^3/\mu\text{L}$ . Range:  $150.00 - 450.00 \times 10^3/\mu\text{L}$ ).
- HIV viral load was undetectable at the last check-up (August 24, 2021).
- The CD4 count was 298 cells per cubic millimeter (August 24, 2021).
- Report of a standard chest radiograph indicates subsegmental atelectasis bibasal, with bilateral obliteration of the costophrenic angles.

A new biopsy (Figure 1), genotyping, and a computed tomography (CT) scan of the abdomen, pelvis, and chest were performed. The histopathology report concluded that it was a giant condyloma acuminata and that there was no evidence of the malignant transformation of the tumor (Figure 1), while the molecular biology report indicated the detection of HPV genotype 11 deoxyribonucleic acid (DNA).

Abdominal and pelvic CT revealed an exophytic lesion at the level of the glans and scrotum throughout its extension and bilateral adenopathies with an infiltrative aspect in the inguinal region (right adenopathy of 10 mm in its greater axis and left of  $16 \times 10$  mm, both accompanied by other smaller ones). Chest CT reported a left lamellar pleural effusion.

Fine-needle aspiration puncture (FNA) of inguinal lymph nodes was performed with the following result: atypical lymphoid cells of undetermined significance, for which the clinical


oncology service recommended performing a biopsy of one of the inguinal lymph nodes. A biopsy of a nodule in the right inguinal region was performed. The following results were obtained: Fibroadipose tissue with chronic inflammation and foreign body-type giant cellular granuloma; no lymph node or neoplastic cells were identified in the sample.

#### Clinical evolution

In the time that elapsed from the initial consultation to performing the biopsy of the inguinal adenopathy, a marked deterioration of the clinical picture was observed, highlighting the local aggressiveness of this entity (Figure 2).

The case was presented at a meeting of a clinical-surgical oncology committee (COCQ), and it was decided not to perform resection surgery due to the size of the lesion, so the patient was referred to the radiotherapy service for definitive treatment. He received 50 Gray (Gy) radiotherapy with a scheme of 1.8 Gy in a single fraction per day from L3 to the distal third of the femur.

Additionally, one month before radiotherapy treatment, treatment with therapeutic vaccination was started using both the systemic and intralesional 9-valent vaccine against HPV (Gardasil 9®). Initially, proposed as a systemic and intralesional guideline, the scheme was at 0, 1, and 4 months.



**Figure 1.** Hematoxylin and eosin-stained skin biopsy – low power. Hyperkeratosis (A), irregular acanthosis with papillomatosis (B), and foci of hypergranulosis with atypical keratinocytes (C) are observed. B: Loss of standard architecture is observed; some keratinocytes present pyknotic nuclei with a whitish halo corresponding to koilocytes (arrows). Some keratinocytes show slight histological pleomorphism.

Due to the size of the lesion, in the initial regimen, the vaccine was injected into the balanopreputial sulcus and into two satellite lesions, diluting 0.5 ml of the vaccine in 2.5 ml of saline solution (SS) and administering between 0.2 and 0.4 ml per injection. Injection site to assess the response to treatment (Figure 3). The patient improved in the clinical picture after the first regimen was decided to use, for the second regimen, 1 ml of the vaccine in 5 ml of saline solution, consistently administering between 0.2 and 0.4 ml per injection site (Figure 4).

At the time that elapsed from the second regimen to the following control (3 months), the lesions recurred on the glans and foreskin, despite having continued treatment with radiotherapy. For this reason, it was decided to increase an intralesional regimen four months after treatment, using 1 ml of a vaccine in 5 ml of SS.

As an adverse effect of the vaccination, he presented pain at the injection site. At the time of writing, 2 of the three systemic regimens (at 0 and 1 months) and 3 of 4 intralesional regimens (0, 1, and 4 months) were administered.

---

## Discussion

TBL is a sexually transmitted disease that appears in 0.1% of the sexually active population and occurs mainly between the fourth and sixth decades of life [ 6 ]. It affects men more frequently (ratio of 2.7:1, some series indicate 4.4:1), but its appearance in women and children has been described in some cases. (7,8) It is very aggressive locally, with a high rate of recurrence (60% - 70%) and malignant potential (30% - 56%) [ 7 ]. The main risk factors for its development are HPV infection, smoking, and immunocompromised states (acquired immunodeficiency syndrome [AIDS], immunosuppressive therapy, alcoholism, diabetes, and chemotherapy) [ 8–10 ].


Infection with HPV subtypes 6 and 11, which are precisely the cause of 90% of condyloma acuminata [ 7, 11 ], is the cause of the disease.

Clinically, it presents as one or several large verrucous lesions with a cauliflower-like appearance that usually ulcerate and may form fistulas [ 7, 12 ]. It generally appears on the external genitalia, although in 10–17%, it can appear in the anorectal region [ 6 ]. Commonly presented symptoms are pain, discomfort, bleeding, the presence of a mass lesion, and discharge. A series of 38 cases reported that the average size and thickness of the tumors reviewed were between 2 – 14 cm and 0.2 – 2.5 cm, respectively, so we can consider that the case we present is rare, as it is more significant than average. [ 8 ].


The diagnosis is mainly clinical, where larger verrucous plaques will be observed than those observed in the case of condylomata acuminata. Histologically, it differs from a simple condyloma in that TBL exhibits increased mitotic activity, significant papillomatosis, thickened tumor margins, and a tendency to penetrate and infiltrate adjacent tissues [ 7 ].

In some series, TBLs that presented more invasion had more frequent keratosis, abnormal mitoses, and microabscesses. Neutrophils represent essential clues to predicting the presence of underlying invasive carcinoma when reviewing biopsy results [ 8 ]. Other essential data to differentiate a giant condyloma acuminata from a squamous cell carcinoma is that the basement membrane is intact in the former, and there is no lymphatic invasion or lymph node metastasis [ 13 ].

Due to the low incidence of this tumor, there are no established guidelines on its treatment yet, but several therapeutic options are described in the literature that must be chosen individually. Treatment currently focuses more on removing verrucous tissues rather than trying to eradicate the virus [ 11 ]. Surgical treatments include surgery with wide margins, and medical treatments such as therapeutic HPV vaccination, imiquimod 5% cream, sinecatechins/polyphenon E, podophyllotoxin, bleomycin, and peptides isolated from amphibians have been described [ 11, 14, 15 ]. Some series suggest that these treatments be followed by physical or chemical ablation if they are large warts [ 16, 17 ].



**Figure 2.** Timeline of the clinical picture.

**Figure**

**3.** Comparative frontals of the clinical picture at 0–2–4–4.1 months. (A) Before starting treatment, (B) one month after administering the second dose of VT and 17 radiotherapy sessions, (C) one month after administering the second dose of VT, and (D) one week after administering a dose of additional intralesional VT. A decrease in the size of satellite lesions (arrows) of the lesions in the glans and balano-preputial sulcus (circle) was observed, but when more than one month passed without VT, the lesions recurred in the glans and foreskin, while the tumor mass that compromised the scrotum remained stable (\*). Circled, arrowed, and asterisked areas injected with the HPV vaccine.



**Figure 4.** Comparative profiles of the clinical picture at 0–1–2–4–4.1 months. (A) Before starting treatment, (B) one month after administering the first dose of the vaccine, (C) one month after administering the second dose of the vaccine and 17 radiotherapy sessions, (D) at four months after starting treatment with VT and ending RT, and (E) at four months, one week after administering an additional intralesional dose of VT. A decrease in the size of satellite lesions (arrows) of the lesions in the glans penis and the balano-preputial sulcus (circle) was observed while the vaccination was monthly, and then the lesions recurred. The decrease in the tumor mass that compromises the scrotum was constant during the four months (\*).

The management of this entity has proven to be complicated, so the focus on identifying risk factors and prevention through HPV vaccination is of the utmost importance. The previously exposed patient presented immunosuppression as a risk factor due to his HIV diagnosis, in addition to being MSM, which could have accelerated the growth of the lesions.

The difficulty of being unable to count on surgery due to tumor extension led to treatment with therapeutic vaccination. In the reviewed literature, only one other article mentions the use of this treatment with an excellent result [15] but taking into account that in that publication, the size of the lesion was significantly smaller (4 cm), in the said study, the bivalent vaccine against HPV systemically, in the present case the nonavalent was used.

Although the therapeutic vaccination was not completed at the time of writing, we noted that there was clinical improvement each time the therapeutic vaccination was administered, even when the additional regimen was administered after the glans and foreskin lesions re-occurred, for more than that the improvement was slight. Regarding radiotherapy, although a slight clinical improvement was observed during treatment, it was not as evident as that observed with vaccination, and once the prescribed regimen was completed, the lesions re-occurred.

In this case, therapeutic vaccination proved helpful and could be considered a first-line treatment in patients with not very advanced tumors or in unresectable tumors to reduce tumor size and evaluate complementary treatment. Aggressive treatment with a more adjusted VT scheme in large tumors seems necessary. TBL is generally associated with specific HPV subtypes (Table 1).

**Table 1.** Correlation of HPV subtypes and their respective diseases [18].

| Illness                   |                                                                                                                                                   | Associated HPV T<br>ype |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------|
| plantar warts             | 1, 2, 4, 63                                                                                                                                       |                         |
| Myrmecia                  | 60                                                                                                                                                |                         |
| common warts              | 1, 2, 4, 26, 27, 29, 41, 57, 65, 77                                                                                                               |                         |
| Common butcher warts      | 1, 2, 3, 4, 5, 10, 28                                                                                                                             |                         |
| flat warts                | 3, 10, 27, 38, 41, 49, 75, 76                                                                                                                     |                         |
| intermediate warts        | 10, 26, 28                                                                                                                                        |                         |
| condyloma acuminata       | 6, 11, 30, 42, 43, 44, 45, 51, 53, 54, 55, 70                                                                                                     |                         |
| Cervical carcinoma        | 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 66, 68, 70                                                                                            |                         |
| laryngeal papillomas      | 6, 11                                                                                                                                             |                         |
| conjunctival papillomas   | 6, 11, 16                                                                                                                                         |                         |
| Buschke–Lowenstein tumor  | 6, 11, 16, 18, 52                                                                                                                                 |                         |
| intraepithelial neoplasms | Low grade: 6, 11, 16, 18, 31, 33, 35, 42, 43, 44, 45, 52, 52, 53, 74<br>High grade: 6, 11, 16, 18, 31, 33, 34, 35, 39, 42, 45, 51, 52, 56, 58, 66 |                         |
| Others                    | 6, 11, 16, 30, 33, 36, 37, 38, 41, 48, 60, 72, 73                                                                                                 |                         |

The vast majority of the literature reviewed suggests that the gold standard is surgery with wide margins [5, 6, 8, 19–23]. Preoperative imaging studies are required to assess the extent of the injury and choose the correct intervention. Despite being the therapy of choice, reinter- ventions are frequently necessary to complete the initial resection due to the high recurrence rate of this tumor [22]. A study with 38 cases reported a recurrence of 23.7% and a mortality of 2.6% at 23 months after undergoing radical surgery [8]. Due to the size of the lesions, their resection can result in extensive defects that may benefit from reconstruction using rotational flaps [1, 2]. Minimally invasive surgery, such as Mohs micrographic surgery, can be used in minor cases, as it has poor results and a high recurrence rate [22].

Regarding the specific treatment in TBL, imiquimod is an immunomodulator that directly activates the cells of innate immunity through Toll-like receptors [7]. It induces the secretion

of proinflammatory cytokines, including interferon-alpha [16]. It also provokes a tremendous inflammatory response by dendritic cells, Langerhans cells, and CD8+ T lymphocytes. It also facilitates the accumulation of CD8+ T lymphocytes in the genital tract, inhibiting tumor growth through gamma interferon. Imiquimod has been shown to induce apoptosis of tumor cells [11]. It is used three times a week for 16 weeks, and local erosions, erythema, and a burning sensation at the application site have been reported as adverse effects [16, 17, 20].

Interleukin 10 inhibitors are also used in treating TBL, a cytokine that exerts an anti-inflammatory action by suppressing the function of macrophages and dendritic cells and acting on T cells, decreasing their production of cytokines. [24]. Imiquimod enhances IL-10 secretion to prevent excessive immune responses and thus prevent autoimmune diseases. Using imiquimod in conjunction with an IL-10 inhibitor has increased efficacy when treating warts [11]. Attention should be given as high or dysregulated levels of IL-10 increase the risk of chronic infection [24].

In some isolated reports, there is treatment with natural peptides isolated from amphibians that are applied in gel form and have been shown to have antimicrobial and anticancer activity. Caerin 1.1 has an anticancer effect against several cancer cell lines in humans (leukemia, lung, colon, melanoma, prostate, ovary, and breast). Caerin 1.9 has broad-spectrum antimicrobial activity and has recently been shown to inhibit the growth of cancer cells in humans. Vitro [11, 16]. It was shown that when both peptides were used in combination with imiquimod, there was strong growth inhibition of HPV+ cells [11]. In addition, both peptides have been shown to have a cytotoxic effect on HPV 16 TC-1 cells and both attract T and NK cells to the tumor site [17].

Administration of the PNGVL4a-CRT/E7 DNA vaccine intralesionally has been shown to elicit a robust immune response by CD8+ T cells in HPV 16+ tumors [25]. In a recent study, a case was treated with systemic 2-valent (Cervarix) and intralesional 9-valent (Gardasil-9) vaccination, with clinical resolution of the condition 12 weeks after starting treatment [15].

Finally, several additional treatments have been described but used in specific situations. Among the local treatments, we found the use of trichloroacetic acid and radiotherapy. However, it can contribute to a malignant transformation of the LBT, for which its use is controversial, the use of CO<sub>2</sub> laser, cryotherapy, and electrofulguration in small cases [7, 26]. Systemic therapy has also been used, among which we find the use of intramuscular or intralesional interferon (consensus interferon, interferon alfa-2a, interferon pegylated) for patients with unresectable tumors in monotherapy or as adjuvant therapy in conjunction with ribavirin, post-operative radiotherapy, 5-fluorouracil, cisplatin, mitomycin, methotrexate, and bleomycin as the most commonly used drugs [6, 7, 11, 26, 27].

Vaccination as a preventive measure is essential since HPV infection is the world's most common sexually transmitted disease [28]. Vaccination of male patients with the quadrivalent HPV vaccine that protects against subtypes 6, 11, 16, and 18 has been shown to significantly reduce the risk of anogenital HPV infection and thus could prevent TBL [7].

---

## Conclusions

Prevention with HPV vaccines in patients with warts is the most critical measure; once the case is advanced, the management of LBT is complex since it is a very aggressive tumor locally, with an easy invasion of neighboring tissues. Resection surgery is the gold standard, and

### Editor's note

Revista Oncología Ecu remains neutral concerning jurisdictional claims on published maps and institutional affiliations.

if it is early in the wrinkle or resectable tumor phase, it is the best therapeutic option. The treatment of cases considered unresectable is complicated, and therapeutic vaccination offers encouraging results as long as the vaccination schedule is adjusted and no more than two months elapse between intralesional doses.

---

## Abbreviations

LBT: Buschke-Lowenstein tumor  
HPV: Human papillomavirus  
HIV: Human Immunodeficiency Virus  
CT: Computed tomography  
DNA: deoxyribonucleic acid  
FNA: fine needle aspiration puncture  
AIDS: Acquired immunodeficiency syndrome.

---

## Administrative information

### Additional Files

The authors declare none.

---

### Acknowledgments

The authors thank all the people of the institutions who collaborated in the development of this research.

---

### Author contributions

1. Conceptualization: Paolo Leone Jiménez, María Fernanda Carvajal.
2. Formal analysis: Paolo Leone Jiménez.
3. Research: Paolo Leone Jiménez, María Fernanda Carvajal.
4. Methodology: Paolo Leone Jiménez.
5. Project administration: María Fernanda Carvajal.
6. Supervision: María Fernanda Carvajal.
7. Validation: Paolo Leone Jiménez.
8. Visualization: Paolo Leone Jiménez, María Fernanda Carvajal.
9. Writing - draft or original: María Fernanda Carvajal.
10. Writing - revision and edition: Paolo Leone Jiménez.

All authors read and approved the final version of the manuscript.

---

### Financing

The authors did not receive any financial recognition for this research work. Expenses incurred in studies and laboratory tests constitute the usual expense of patients admitted to the institution and did not constitute an additional expense to patients. The authors subsidized the administrative costs of this research.

---

### Availability of data and materials

Data availability is available upon request to the corresponding author. No other materials were reported.

---

## Statements

### Ethics committee approval

It does not apply to observational studies with a review of databases or medical records.

---

### Consent to publication

The authors have permission for publication from the patient referred to in this case.

---

### Conflicts of interest

The authors declare that they have no conflict of interest or competence.

---

## References

1. Steffen C. The men behind the eponym--Abraham Buschke and Ludwig Lowenstein: giant condyloma (Buschke-Loewenstein). *Am J Dermatopathol*. 2006 Dec;28(6):526-36. doi: [10.1097](#). PM ID: [17122499](#).
2. Nordsiek M, Ross C, Metro M. Successful Surgical Management of Giant Condyloma Acuminatum (Buschke Lowenstein Tumor) in the Urethra of a Female Patient: A Case Report. *Curr Urol* 2015 May;8(1):49- 52. DOI: 10.1159/000365689. Epub 2015 May 20. PM ID: [26195964](#); PM CID: PMC4483283.
3. Valenzuela N, Barnadas MA, Matias-Guiu X, Curell R, de Andres L, Alomar A. Lethal perianal Buschke-Löwenstein tumor in a female patient with AIDS. *Dermatology*. 2004;208(4):349 -50. doi: 10.1159/000077847. PM ID: [15178921](#).
4. Chu QD, Vezeridis MP, Libbey NP, Wanebo HJ. Giant condyloma acuminatum (Buschke-Lowenstein tumor) of the anorectal and perianal regions. Analysis of 42 cases. *Dis Colon Rectum*. 1994Sep;37(9):950-7. DOI: 10.1007/BF02052606. PM ID: [8076499](#).
5. Fai EK, Bhutta HA, Ali K. Progression of a Buschke-Lowenstein tumor to invasive squamous cell carcinoma. *JAAPA: Official Journal of the American Academy of Physician Assistants*. 2019 Jun;32(6):1-4. DOI: 10.1097/01.jaa.0000554740.24182.ae. PM ID: [31136407](#).
6. Ben Kridis W, Werda I, Charfi S, Toumi N, Boudawara T, Mzali R, Daoud J, Khanfir A. Buschke - Lowenstein anal tumor: an ambiguous entity. *Exp Oncol*. 2019 Jun;41(2):182-184. DOI: 10.32471/exp-oncology.2312-8852.vol-41-no-2.13119. PM ID: [31262155](#).
7. Nieves-Condoy JF, Acuña-Pinzón CL, Chavarria-Chavira JL, Hinojosa-Ugarte D, Zúñiga-Vázquez LA. Giant Condyloma Acuminata (Buschke-Lowenstein Tumor): Review of an Unusual Disease and Difficult to

- Manage. Infect Dis Obstet Gynecol. 2021 Jun 30;2021:9919446. doi: 10.1155/2021/9919446. PMID:[34305393](#); PMCID: PMC8266468.
8. Zhang D, Gonzalez RS, Feely M, Umrau K, Lee H, Allende DS, Karamchandani DM, Zaleski M, Lin J, Westerhoff M, Zhang X, Alpert L, Liao X, Lai J, Liu X. Clinicopathologic features of Buschke - Löwenstein tumor: a multi-institutional analysis of 38 cases. *Virchows Arch*. 2020 Apr;476(4):543-550. doi:10.1007/s00428-019-02680-z. Epub 2019 Nov 15. PMID: [31728626](#).
  9. De Araújo PSR, Padilha CEG, Soares MF. Buschke-Lowenstein tumor in a woman living with HIV/AIDS. Vol. 50, *Journal of the Brazilian Society of Tropical Medicine*. Brazilian Society of Tropical Medicine; 2017. p. 577. S U: [www.scielo.br](http://www.scielo.br)
  10. Wiedemann A, Diekmann WP, Holtmann G, Kracht H. Report of a case with giant condyloma (Buschke-Löwenstein tumor) localized in the bladder. *J Urol*. 1995 Apr;153(4):1222-4. PMID: [7869506](#).
  11. Yuan J, Ni G, Wang T, Mounsey K, Cavezza S, Pan X, Liu X. Genital warts treatment: Beyond imiquimod. *Hum Vaccin Immunother*. 2018 July 3;14(7):1815-1819. DOI: 10.1080/21645515.2018.1445947. Epub 2018 April 9. PMID: [29505317](#); PMCID: PMC6067868.
  12. Lee PK, Wilkins KB. Condyloma and other infections, including human immunodeficiency virus. *Surg Clin North Am*. 2010 Feb;90(1):99-112, Table of Contents. DOI: 10.1016/j.suc.2009.09.005. PMID: [20109635](#).
  13. Knoblich R, Failing JF Jr. Giant condyloma acuminatum (Buschke-Löwenstein tumor) of the rectum. *Am J Clin Pathol*. 1967 Oct;48(4):389-95. doi: 10.1093/ajcp/48.4.389. PMID: [6061519](#).
  14. Martin JM, Molina I, Monteagudo C, Marti N, Lopez V, Jorda E. Buschke-Lowenstein tumor. *J Dermatol Case Rep*. 2008 Dec 27;2(4):60-2. DOI: 10.3315/jdcr.2008.1019. PMID: [21886716](#); PMCID: PMC3157782.
  15. Thomas R, Smith-Matthews S, Ho J. Giant condyloma of Buschke-Lowenstein in a patient with pemphigus vegetans treated with intralesional and systemic human papillomavirus vaccine. *JAAD Case Rep*. 2021 Dec 29;20:54-57. DOI:10.1016/j.jdcr.2021.12.024. PM ID: [35071717](#); PM CID: PMC8761607.
  16. Ma B, Yuan J, Chen S, Huang K, Wang Q, Ma J, Lin R, Zhang L, Zhou Y, Wang T, Walton SF, Pan X, Chen G, Wang Y, Ni G, Liu X. Topical application of temperature-sensitive caerin 1.1 and 1.9 gel inhibits TC-1 tumor growth in mice. *Am J Transl Res*. 2020 January 15;12(1):191-202. PMID: [32051748](#); PMCID: PMC7013226.
  17. Chen S, Zhang P, Xiao L, Liu Y, Wu K, Ni G, Li H, Wang T, Wu X, Chen G, Liu X. Caerin 1.1 and 1.9 Peptides from Australian Tree Frog Inhibit Antibiotic-Resistant Bacteria Growth in a Murine Skin Infection Model. *Microbiol Spectr*. 2021 Sep 3;9(1):e0005121. DOI: 10.1128/Spectrum.00051-21. Epub 2021 July 14. PMID: [34259550](#); PMCID: PMC8552723.
  18. Klaus Wolff, Richard Allen Johnson, Arturo P. Saavedra EKR. Fitzpatrick's Color Atlas and Synopsis of Clinical Dermatology. Eight Edition. 2017. 1992 p. S U: [accessmedicine.mhmedical.com](http://accessmedicine.mhmedical.com)
  19. Ibrahim A, Ziani I, correspondent A. Cite this article: Ahmed Ibrahim et al. Tumeur de Buschke-Lowenstein à localisation pénienne. *Pan African Medical Journal*. 2020;37:37. SU: [www.panafrican-med-journal.com](http://www.panafrican-med-journal.com)
  20. Sonthalia S, Gandhi V, Agrawal M, Sharma P. Successful nonsurgical treatment of penile Buschke-Löwenstein tumor with 12 weeks of 5% imiquimod alone. *Int J STD AIDS*. 2019 Oct;30(11):1136-1139. DOI: 10.1177/0956462419855505. Epub 2019 September 24. PMID: [31551001](#).
  21. Bertram P, Treutner KH, Rübber A, Hauptmann S, Schumpelick V. Invasives Schuppenzellkarzinom beim Riesenanorektalkondylom (Buschke-Löwenstein-Tumor). *Langenbecks Archiv für Chirurgie*. 1995 Mar;380(2):115-8. S U: [scielo.conicyt.cl](http://scielo.conicyt.cl)

22. Spinu D, Rădulescu A, Bratu O, Checheriță IA, Ranetti AE, Mischianu D; -. Giant condyloma acuminatum - Buschke-Lowenstein disease - a literature review. Surgery (Bucur). 2014 Jul-Aug;109(4):445-50. PMID: [25149605](#).
23. Kadouri Y, Nouini Y. La tumeur de Buschke-Löwenstein [Buschke-Löwenstein' tumor]. Pan Afr Med J. 2020 Aug 28;36:359. French. doi: 10.11604/pamj.2020.36.359.13292. PMID: [33224425](#) ; PMCID: PMC7664135.
24. O'Garra A, Barrat FJ, Castro AG, Vicari A, Hawrylowicz C. Strategies for the use of IL-10 or its antagonists in human disease. Immunol Rev. 2008 Jun;223:114-31. DOI: 10.1111/j.1600-065X.2008.00635.x. PMID: [18613832](#).
25. Alvarez RD, Huh WK, Bae S, Lamb LS Jr, Conner MG, Boyer J, Wang C, Hung CF, Sauter E, Paradis M, Adams EA, Hester S, Jackson BE, Wu TC, Trimble CL. A pilot study of pNGVL4a-CRT/E7(detox) for the treatment of patients with HPV16+ cervical intraepithelial neoplasia 2/3 (CIN2/3). Gynecol Oncol. 2016 Feb;140(2):245-52. DOI: 10.1016/j.ygyno.2015.11.026. Epub 2015 November 23. PMID: [26616223](#) ; PMCID: PMC4724445.
26. Valle, Adriana Della, Michel Heguaburu, Rodrigo Fresco, Gloria Roldan, and Monica Della Valle. "Buschke- Löwenstein tumor: case report and review with focus on therapeutic and psychological aspects." Internet Journal of Surgery 9 (2006): no. p. S U: [www.semanticscholar.org](http://www.semanticscholar.org)
27. Shrestha NK, Hamrock DJ. Successful treatment of disseminated human papillomavirus infection with pegylated interferon and ribavirin. Clin Infect Dis. 2010 July 1;51(1):e4-6. DOI: 10.1086/653428. PMID: [20504235](#).
28. Pennycook KB, McCready TA. Condyloma Acuminata. 2022 May 8. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. PM ID: [31613447](#).
29. Kang S, Amagai M, Bruckner AL, Enk AH, Margolis DJ, McMichael AJ, et al. Fitzpatrick's Dermatology. 9th Edition. McGraw-Hill Education; 2019. S U: [accessmedicine.mhmedical.com](http://accessmedicine.mhmedical.com)