

Hyperwarburgism: the "happy hypoglycemia" of cancer patients: Narrative review

*Correspondence:

Fernando.foosemberg1@hotmail.com

Address:

Av. Pedro J. Menéndez Gilbeir y Atahualpa Chávez (junto a la ciudadela Atahazana)
Apt. Postal 090505
Unidad de Cuidados Intensivos.
Instituto Oncológico Nacional "Dí. Juan I'anca Maiengo" SOLCA
Guayaquil - Ecuador. Phone: +593-43718300 EXI. 2218

Conflict of interest: The authors declare to have no conflicts of interest.

Received: August 15, 2022

Accepted: October 26, 2022

Published: December 7, 2022

Editor: Dr. Evelyn Valencia Espinoza.

Cite:

Roosemberg F, Veliz J, Ortega V, Bowen A, García C. Hyperwarburgism: the "happy hypoglycemia" of cancer patients: Narrative review. Rev. Oncol. Ecu 2022;32(3):334-342.

ISSN: 2661-6653

DOI: <https://doi.org/10.33821/626>



Copyright Roosemberg F, et al. This article is distributed under the terms of [Creative Commons Attribution License BY-NC-SA 4.0](#), which allows the use and redistribution citing the source and the original author.

Fernando Roosemberg Ordoñez ¹ , **José Veliz Ortega** ¹ , **Valeria Ortega Uscocovich** ¹ , **Aquiles Bowen Flores** ¹ , **Carlos García Cruz** ¹

1. Intensive Care Unit, SOLCA - Guayaquil.

Summary

Introduction: Glucose is the cerebral energy fuel; this relationship is fully established in most reviews due to neuronal tissue's avid -and almost exclusive- glucose consumption. In this sphere, hypoglycemia is translated by default into a set of neurological symptoms resulting from the neuroglycopenic state. When the drop in these glycemic levels is pronounced, it triggers alterations in the sensory state, being able to reach a coma with irreversible damage if sustained over time.

Purpose of the review: The objective is to present a case of severe hypoglycemia without neuroglycopenic symptoms.

Recent findings: The absence of neurological symptoms is due to the consumption of Lactate – traditionally an anaerobic product – as an alternative energy metabolic pathway to glucose consumption. Hypoglycemia can be compensated at the neurological level with lactate launching systems in neuronal tissue, replacing glucose as the brain's energy substrate.

Conclusions: Hypoglycemia without adrenergic or neuroglycopenic symptoms is an issue linked to cancer patients, and proposes Lactate as fuel for nervous tissue in addition to glucose. On the other hand, the Lactate = hypoperfusion association is another entity that must be reviewed and reanalyzed for everything that Lactate implies within the body's metabolic pathophysiological pathway.

Keywords :

DeCS: Hypoglycemia, Lactic Acid, Pyruvic Acid, Medical Oncology, Anaerobiosis, Metabolic Encephalopathies.

Abstract

Introduction:

MESH: Hypoglycemia; Lactic Acid; Pyruvic Acid; Medical Oncology; Anaerobiosis; Brain Diseases, Metabolic.

Introduction

The complexity of the known metabolic pathways can only be overshadowed by the number of alternatives to traditional pathways that could be developed at the cellular level, many of which have been studied and others whose physiological or pathophysiological explanation is not well known.

Within this context, the oncology section deserves particular mention. Neoplastic cells are characterized by a unique metabolism, developed with specific characteristics to replace a high energy demand necessary for excessive proliferation. The goal of the tumor is to divide and grow; for this to happen, it must be in a hypoxic environment [1] and with a high lactate concentration [2].

Each tumor to fulfill its pathological proliferation is governed by a phenomenon described in 1920 by Otto Warburg called the " Warburg effect " [3]. This effect is shown as an alteration of the normal metabolism of glucose due to the exaggerated increase in its consumption by neoplastic cells. However, unlike normal cells, cancer cells take an alternative route to produce energy; the glucose, instead of being transported to the mitochondria, is transported to the cytoplasm, and that is where Lactate is transformed into pyruvate to access the tricarboxylic acid pathway (a process mediated through the enzyme lactate dehydrogenase) and is then secreted (aerobic glycolysis) [3].

Glycolysis causes a change in the environment of the tumor cell area. Elevated Lactate in the body acts as a cellular energy substrate, which protects against hypoglycemic events [4]

. As the study of Lactate progresses, this " Warburg effect " is not exclusive to cancer cells, as it can also occur in other tissues. The effect of not presenting adrenergic signs or symptoms or neuroglycopenia has been reported in cancer patients presenting episodes of hypoglycemia since Lactate also feeds neuronal cells, a rare process described as hyperwarburgism [5]. Little literature on hyperwarburgism is described in the leading medical reference libraries, the vast majority of cancer patients (66% of cases).

Materials and methods

Study design

A search was performed in the United States National Library of Medicine (MEDLINE) data- base through The United States National Library of Medicine and National Institutes of Health (PUBMED) with the following search details:

- Date: between the years 1960 and 2021
- Languages: all those of the search engine
- Terms used: using the terms " hypoglycemia " + " warburg " + " effect ".

Nineteen articles were identified, including bibliographic reviews, analyses, and investigations, along with three review articles of 3 cases, 2 of which were linked to neoplastic processes and 1 in a non-oncological patient. It is mentioned that other keywords were used without variability in the results obtained, and the search tree by MESH Terms had no different consequences.

Definitions

The precise definition of hypoglycemia is currently under debate; however, the American Diabetes Association (ADA) describes hypoglycemia as a clinical condition characterized by low blood glucose concentrations, usually less than 70 mg/dL (3.89 mmol/L) [6]. Diagnosis of hypoglycemia is based on Whipple's triad, which is described as low blood glucose, symptoms of hypoglycemia, and symptom improvement once glycemia normalizes [7]. We can classify the symptoms of hypoglycemia as: adrenergic (tachycardia, palpitations, tremors, sweating, paleness, and anxiety) and non-adrenergic or neuroglycopenic (hunger, headache, weakness, visual disturbances, confusion, lethargy, seizures, and even coma) [7]. A differential diagnosis would be considered (due to the presence of hypoglycemia without etiology) with insulinoma, which was not captured by the imaging studies carried out within its oncological analysis (para-clinical rule out) together with the absence of neuroglycopenic symptoms as cited (clinical rule out). Glucose is the primary substrate used by the brain. As such, multiple regulatory mechanisms exist to maintain blood glucose concentrations within normal parameters [8]; however, it is not the only substrate from which the brain can derive energy [9]. Showing Lactate as a viable energy alternative. Experimental and clinical data provide many arguments in favor of this hypothesis, mainly suggesting that Lactate is a possible and sometimes even mandatory substrate for the brain in an energy crisis [9]. Lactate, from 1780 and even up to recent years, was analyzed and described as the final waste product of metabolic pathways under anaerobic conditions with no other pathway than its subsequent clearance, formed in times of poor perfusion or hypoxia, an important marker of organ dysfunction [10] and its usefulness outside of being an indicator of the processes above of low perfusion and sepsis has been questioned for decades [11].

Lactate in the literature

Reviewing the literature on Lactate, like other authors, we will mention in the first place one of the body's energy deposit sites: skeletal muscle. We know that skeletal muscle is a storage fuel, being a reservoir of glycogen that can be broken down to Lactate (not just glucose [12]) in response to the binding of β -adrenergic stimulants to the muscle cell membrane [12]. Lactate generated after beta-adrenergic stimulation is distributed passively down the concentration gradient out of the cell, passing into other tissues via monocarboxylate transporters (MCTs), where it can be used as fuel [4, 14]. In his research on Lactate, Daniel specifies that this mechanism in traumatic situations, stress, or high energy requirements, allows rapid fuel consumption at low cost [13]. Lactic acid dehydrogenase, or lactate dehydrogenase (LDH), is an essential enzyme of the anaerobic metabolic pathway that catalyzes the reaction of pyruvate to Lactate [14]. Lactate dehydrogenase has higher activity than other enzymes in glycolysis or oxidative metabolism [14]. As a result of LDH activity, Lactate is constantly formed and consumed almost instantaneously [15], which is why its serum levels under normal conditions are incipient (<2mmol/L).

Astrocyte lactate transporter system (ANLS)

Since the body has a large deposit in glycogen, Lactate, in particular situations, becomes an excellent source of fuel for most tissues, and the central nervous system, which our review includes, is no exception. Pellerin and Magistretti 1994 introduced the astrocyte-neuron lactate transporter (ANLS). This transporter system is proposed to molecularly explain the consumption of Lactate instead of glucose in neuronal tissue. Under the premises of this system,

astrocytes take up glucose (influenced by glutamate re-uptake) and then release Lactate for use by neurons attached to the system [15].

In ANLS, the astrocyte makes the energetic link between the capillary and the neuron through the intercellular lactate transport system. When electrical stimulation produces neuronal activation (neuronal membrane activity), large amounts of glutamate (among other neurotransmitters) are released into the synaptic cleft [16]. This glutamate neurotransmitter must be quickly reabsorbed to be transformed into glutamine, which requires energy. Glutamate re-uptake in astrocytes triggers increased glucose uptake from capillaries (an energy-supplying molecule), a process regulated by isoform 2 of the Na⁺/K⁺-ATPase pump [17]. The generation of Lactate provides the energy required by the astrocyte for these processes (ATP) from actively captured capillary glucose (glycolysis), a process regulated by Lactate Dehydrogenase (LDH) isoenzyme 5 [18]. Lactate produced within the astrocyte must continue to the neuron. This lactate exchange is carried out thanks to the monocarboxylate transporters 1 and 4 (MCT1 and MCT4), which according to the approach, are present in the astrocyte [19]. Bittar et al., in their immunohistochemical and positron emission tomography (PET) studies, support the notion of a regulated lactate flow between astrocytes and neurons with the participation of the different LDH isoenzymes [20]. In this way, the Lactate produced inside the astrocyte mediated by the LDH5 isoenzyme will be recaptured by the neurons through the monocarboxylate transporter 2 (MCT2), which is present in the neuron, to join the Krebs cycle, prior to the action of LDH isoenzyme one that converts Lactate into pyruvate, now as pyruvate to be part of the tricarboxylic acid pathway, resulting in obtaining energy for the maintenance of neuronal functions [20] and continuing synaptic transmission with the production of neurotransmitters glutamate, aspartate and gamma-aminobutyric acid (GABA) [17, 18, 19] glutamate that will return to the presynaptic space and will be re-uptake by the astrocyte, to repeat the cycle (Figure 1).




Figure 1. Schematic representation of proposed lactate transport between astrocytes and neurons (ANLS). (Pellerin and Magistretti, 1994, 1997 – Modified by authors). Capillary glucose performs via glycolysis the production of pyruvate, which, in the presence of high concentrations of LDH5 without the need for anaerobiosis, is transformed into Lactate, which passes to the neuron via MCT 1 and 2 to return to its tricarboxylic form of pyruvate mediated by LDH1 and participate in the Krebs cycle.

MCT analysis

MCTs are a proper input/output system that allows molecular flow between plasma and neuronal tissue. That Lactate is continuously formed and used under fully aerobic conditions has been a proven truth for decades [12]. The discoveries of many lactate exchangers at the tissue level, including with different cell lines, have led to the articulation of numerous "lactate transporter systems" between producer and consumer cells [18]. The Pellerin / Magistretti model proposes MCT1 and MCT4 as those responsible for lactate export by astrocytes. In contrast, MCT2 has been proposed as a transporter that incorporates Lactate into neurons [15, 19], leading to thus the flow of Lactate between both cells for the final use of Lactate in the Krebs cycle [20]. This proposed hypothesis fits with the operation of the ANLS system. However, Ferguson, in his study, considers that all MCTs are not unidirectional; that is, they allow the flow of the substrate in both directions [1 0]; studies on their distribution have shown that their expression is specific in certain tissues: MCT1 is expressed in astrocytes, oligodendrocytes, endothelial cells, and many other tissues throughout the body, MCT4 is expressed in astrocytes, but not in neurons (as well as in muscle cells and some other places in the body), and MCT2 are found predominantly in neurons [15, 19]. Although the concept of ANLS and the experiments that support it have been debated, the importance of Lactate within the central nervous system (CNS) is evident. It is a valid hypothesis to justify clinical processes in hypoglycemic/hyperlactatemia states.

Hyperwarburgism in the medical literature

3 reports of hyperwarburgism were found in the literature. The case presented by Elhosmy et al. corresponds to a 64-year-old woman who, despite presenting hypoglycemia, did not have any clinical manifestation of neuroglycopenia, despite serum glucose of 26 mg/dL (1.4 mmol /liter) and serum lactate of 28.5. mmol /L. The patient presented with an oncological diagnosis of diffuse large B-cell lymphoma. Despite the dextrose infusions, they reported no increase in blood glucose, but paradoxically there was an increase in serum lactate. Goyal et al. present the case of a 52-year-old man with severe but asymptomatic hypoglycemia in the oncologic setting of diffuse large B-cell lymphoma [21]. Loeb et al., however, presented a non-cancer patient: an 85-year-old woman who presented with hyperwarburgism in the context of hemorrhagic shock with no cancer history [22].

Table 3. Hyperwarburgism in the medical literature.

Authors	Patient	Oncology	hypoglycemia	hyperlactatemia	neuroglycopenia
Elhosmy et al	female, 64a	Yes: Diffuse B-cell lymphoma	Yes	Yes	Nope
goyal et al	more, 52a	Yes: Diffuse B-cell lymphoma	Yes	Yes	Nope
Loeb et al	female, 85th	Nope	Yes	Yes	Nope

The ANLS system proposed by Pellerin and Magistretti satisfactorily provides the molecular explanation of the process: in states of hypoglycemia with decreased glucose supply to the neuronal tissue, this, via the neuronal astrocyte lactate transporter, takes up Lactate produced in the astrocyte to maintain the energy supply of the neuron in order to continue its functions. This process is one of the many that continues to accumulate data in favor of Lactate as the leading participant in metabolism and that neurons have alternative pathways to obtain their

energy, not only in cancer patients (where there is more evidence) but also in non-cancer patients where studies would be lacking to demonstrate the case of Loeb.

conclusions

Hypoglycemia without adrenergic or neuroglycopenic symptoms is a rare topic in literature reviews. Widely linked to cancer patients but not exclusive to them, it is no longer a metabolic curiosity to become a valid topic of interest in neuronal metabolism since the sustained belief that glucose is the only fuel for tissue is in check highly strung. The analysis would remain to determine under what circumstances lactate comes into play through the ANLS to generate energy since there are cases in which it occurs outside the neoplastic environment. Knowing and elucidating the mechanisms that condition the use of alternative neural energy pathways would allow a better understanding of brain metabolism, which would open doors to search for protective therapies in cases of hypoperfusion or drop in energy intake for different reasons and would allow us to provide adequate support to care for the neural system. On the other hand, the Lactate = hypoperfusion association is another entity that must be reviewed and reanalyzed for everything that Lactate implies within the body's metabolic pathophysiological pathway.

Acknowledgments

Not applicable.

Administrative information

Abbreviations

ANLS: Neuronal astrocyte lactate transporter system.

MCT: monocarboxylate transporter.

CNS: central nervous system.

Additional Files

None declared by the authors.

Financing

The authors did not receive any financial recognition for this research work. The expenses incurred in this study constitute the regular activity of the oncology service, so they did not constitute an additional expense for the participants.

Availability of data and materials

Data is available upon request to the corresponding author. No other materials are reported.

Author contributions

Fernando Roosenberg Ordoñez: Conceptualization, Formal analysis, Research, methodology, Project administration, Supervision, Validation, Visualization, Writing-original draft, Writing-revision, and editing.

José Veliz Ortega: Conceptualization, formal analysis, research, validation, writing-revision, and editing.

Valeria Ortega Usocovich: Conceptualization, formal analysis, research, validation, writing-revision, and editing.

Aquiles Bowen Flores: Supervision, validation, writing-revision, and editing:

Carlos García Cruz: formal analysis, research, methodology, validation, visualization, writing -original draft.

All authors read and approved the final version of the manuscript.

Ethics committee approval

It does not apply to observational studies and narrative reviews.

Consent for publication

It does not apply when the patients' images, x-rays, or tomographies are not published.

References

1. Arvelo F, Cotte C. Hypoxia in cancer malignancy: Review. *Investig Clin*. 2009;50 (4):529-546. SCIELO: [SKY](#): S0535.
2. Rengifo GFG, Castañeda CG, Guerinoni DE, Tubeh R. Gene expression of glycolytic pathway enzymes in cancer cells. Published online 2007:11. SCIELO: [SKY](#): S1728.
3. Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg Effect: The Metabolic Requirements of Cell Proliferation. *Science*. 2009;324(5930):1029-1033. DOI: [10.1126/science.1160809](#) PMid:19460998 PMID: PMC2849637
4. Goodwin ML, Gladden LB, Nijsten MWN. Lactate-Protected Hypoglycemia (LPH). *Front Neurosci*. 2020;14:920 . DOI: [10.3389/fnins.2020.00920](#) PMid: 33013305 PMID: PMC7497796
5. Elhomisy GC, Eranki V, Albert SG, et al. "Hyper - Warburgism," a Cause of Asymptomatic Hypoglycemia with Lactic Acidosis in a Patient with Non-Hodgkin's Lymphoma. *J Clin Endocrinol Metab* . 2012;97(12):4311- 4316. DOI: [10.1210/jc.2012-2327](#) PMid: 23055548
6. Nares-Torices MÁ, González-Martínez A, Martínez-Ayuso FA, et al. hypoglycemia: time is the brain. What are we doing wrong? *Mid Int Mex*. 2018;34(6):881-895. DOI: [10.24245/mim.v34i6.2040](#)

7. Gutierrez Medina S, Aragon Valera C, Dominguez Fernandez R, Garcia Sanchez L, Manrique Franco K, Rovira Loscos A. Factitious hypoglycemia. *Endocrinol Nour.* 2013;60(3):147-149. DOI: [10.1016/j.endonu.2012.01.024](https://doi.org/10.1016/j.endonu.2012.01.024) PMid: 22591994
8. Kittah NE, Vella A. Management of endocrine disease: Pathogenesis and management of hypoglycemia. *Eur J Endocrinol.* 2017;177(1):R37-R47. DOI: [10.1530/EJE-16-1062](https://doi.org/10.1530/EJE-16-1062) PMid: [28381450](https://pubmed.ncbi.nlm.nih.gov/28381450/)
9. Orban JC, Leverve X, Ichai C. Lactate: the energy substrate of tomorrow. *Intensive care.* 2010;19 (5):384 -392. DOI: [10.1016/j.reaurg.2010.05.016](https://doi.org/10.1016/j.reaurg.2010.05.016)
10. Ferguson BS, Rogatzki MJ, Goodwin ML, Kane DA, Rightmire Z, Gladden LB. Lactate metabolism: historical context, prior misinterpretations, and current understanding. *Eur J Appl Physiol.* 2018;118 (4):691-728. DOI: [10.1007/s00421-017-3795-6](https://doi.org/10.1007/s00421-017-3795-6) PMid: 29322250
11. Owen OE, Morgan AP, Kemp HG, Sullivan JM, Herrera MG, Cahill GF. Brain Metabolism during Fasting*. *J Clin Invest.* 1967;46(10):1589-1595. DOI: [10.1172/JCI105650](https://doi.org/10.1172/JCI105650) PMid: 6061736 PMID: PMC292907
12. Brooks GA. Lactate shuttles in nature. *Biochem Soc Trans.* 2002 Apr;30 (2):258-64. PMID: 12023861. PMID: [12023861](https://pubmed.ncbi.nlm.nih.gov/12023861/) DOI: [10.1042/bst0300258](https://doi.org/10.1042/bst0300258)
13. Daniel AM, Shizgal HM, MacLean LD. The anatomical and metabolic source of Lactate in shock. *Surge Gyne-col OB.* 1978 Nov;147 (5):697-700. PMID: [715647/](https://pubmed.ncbi.nlm.nih.gov/715647/)
14. Farhana A, Lappin SL. Biochemistry, Lactate Dehydrogenase. 2021 May 7. In: StatPearls [Internet]. TreasureIsland (FL): StatPearls Publishing; 2022 Jan-. PMID: [32491468/](https://pubmed.ncbi.nlm.nih.gov/32491468/)
15. Brooks GA. The Science and Translation of Lactate Shuttle Theory. *Cell Metab.* 2018;27 (4):757-785. DOI: [10.1016/j.cmet.2018.03.008](https://doi.org/10.1016/j.cmet.2018.03.008) PMid: 29617642
16. Pellerin L, Magistretti PJ. Excitatory amino acids stimulate aerobic glycolysis in astrocytes via an activation of the Na⁺/K⁺ ATPase. *Dev Neurosci.* 1996 ;18(5-6):336-342. DOI: [10.1159/000111426](https://doi.org/10.1159/000111426) PMid 8940604
17. Pellerin L, Magistretti PJ. Glutamate uptake stimulates Na⁺, K⁺-ATPase activity in astrocytes via activation of a distinct subunit highly sensitive to ouabain. *J Neurochem .* 1997 ;69 (5):2132-2137. DOI: [10.1046/j.1471-4159.1997.69052132.x](https://doi.org/10.1046/j.1471-4159.1997.69052132.x) PMid: 9349559
18. Pellerin L, Pellegrini G, Bittar PG, et al. Evidence Supporting the Existence of an Activity-Dependent Astrocyte- Neuron Lactate Shuttle. *Dev Neurosci.* 1998;20(4-5):291-299. DOI: [10.1159/000017324](https://doi.org/10.1159/000017324) PMid: 9778565
19. Pellerin L, Pellegrini G, Martin JL, Magistretti PJ. Expression of monocarboxylate transporter mRNAs in mouse brain: Support for a distinct role of Lactate as an energy substrate for the neonatal vs. adult brain. *Proc Natl Acad Sci.* 1998;95(7):3990-3995. DOI: [10.1073/pnas.95.7.3990](https://doi.org/10.1073/pnas.95.7.3990) PMid: 9520480 PMID: PMC19950
20. Bittar PG, Charnay Y, Pellerin L, Bouras C, Magistretti PJ. Selective distribution of lactate dehydrogenase iso- enzymes in neurons and astrocytes of the human brain. *J Cereb Blood Flow Metab Off J Int Soc Cerebral Blood Flow Metab.* 1996;16 (6):1079-1089. DOI: [10.1097/00004647-199611000-00001](https://doi.org/10.1097/00004647-199611000-00001) PMid:8898679
21. Goyal I, Ogbuah C, Chaudhuri A, Quinn T, Sharma R. Confirmed Hypoglycemia Without Whipple Triad: A Rare Case of Hyper-Warburgism . *J Endocr Soc.* 2021;5(1):bvaa182. DOI: [10.1210/ jendso /bvaa182](https://doi.org/10.1210/ jendso /bvaa182) PMid: 33354638 PMID: PMC7737393
22. Loeb T, Ozguler A, Baer G, Baer M. The pathophysiology of "happy" hypoglycemia. *Int J Emerg Med.* 2021;14(1):23. DOI: [10.1186/s12245-021-00348-7](https://doi.org/10.1186/s12245-021-00348-7) PMid: 33882828 PMID: PMC8058752

Editor's Note

Revista Oncología Ecu remains neutral concerning jurisdictional claims in published maps and institutional affiliations.